

## 2009 MQIC MEASURES SUMMARY LISTING FOR ANNUAL PERFORMANCE REPORTING

### DIABETES

#### HEDIS® Measures:

##### 1. Comprehensive Diabetes Care (CDC):

Percentage of members 18-75 years of age with diabetes (type 1 and type 2) who had each of the following:

- a. HbA1c Testing
- b. HbA1c Poor Control (>9.0%)
- c. HbA1c Control (<8.0%)
- d. HbA1c Good Control (<7.0%) (Additional exclusion criteria are required for this indicator)
- e. Eye Exam (retinal) performed
- f. LDL-C Screening
- g. LDL-C Control (<100 mg/dL)
- h. Medical Attention for Nephropathy
- i. Blood Pressure Control (<130/80 mm Hg)
- j. Blood Pressure Control (<140/90 mm Hg)

#### Non-HEDIS® Measures:

##### 1. Statin Therapy for Persons with Diabetes

Members with at least one statin prescription filled during the measurement year.

##### 2. Disease-Burden Measures (reported every 5 years; due with HEDIS 2013 data)

- a. Acute Myocardial Infarction (AMI)
- b. Cerebrovascular Accident (CVA)
- c. Amputations
- d. Dialysis

### ASTHMA

#### HEDIS® Measures:

##### 1. Use of Appropriate Medications for People With Asthma (ASM):

Percent members 5-56 years of age who were identified as having persistent asthma and who were appropriately prescribed medication.

- a. 5-9 Years Old
- b. 10-17 Years Old
- c. 18-56 Years Old
- d. Total (sum of the three numerators divided by sum of the three denominators)

#### Non-HEDIS® Measures:

##### 1. Periodic Assessment (5-56 Years Old)

Members who have had at least one preventive visit during the measurement year.

##### 2. Emergency Department Visits for Members with Asthma and Being Treated for a Primary Diagnosis of Asthma

## **TOBACCO CONTROL**

### **HEDIS<sup>®</sup> Measures:**

#### **1. Medical Assistance With Smoking Cessation (MSC):**

The following components assess different facets of providing medical assistance with smoking cessation:

- a. Advising Smokers to Quit
- b. Discussing Smoking Cessation Medications
- c. Discussing Smoking Cessation Strategies

### **Non-HEDIS<sup>®</sup> Measure:**

#### **1. Percent Current Smokers**

-----

## **DEPRESSION**

### **HEDIS<sup>®</sup> Measures:**

#### **1. Antidepressant Medication Management (AMM):**

Percent members 18 years of age and older who were diagnosed with a new episode of major depression, treated with antidepressant medication, and who remained on an antidepressant medication treatment.

- a. Effective Acute Phase Treatment
  - b. Effective Continuation Phase Treatment
- 

## **SUBSTANCE USE DISORDERS**

### **HEDIS<sup>®</sup> Measures:**

#### **1. Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET):**

Percent of adolescent and adult members with a new episode of alcohol or other drug (AOD) dependence who received the following:

- a. Initiation of AOD Treatment
  - b. Engagement of AOD Treatment
- 

## **HYPERCHOLESTEROLEMIA**

### **HEDIS<sup>®</sup> Measures:**

#### **1. Cholesterol Management for Patients with Cardiovascular Conditions (CMC):**

Percent members 18-75 years of age who were discharged alive for AMI, CABG or PTCA from January 1 to November 1 of the year prior to the measurement year, or who had a diagnosis of IVD during the measurement year and the year prior to measurement year, who had each of the following during the measurement year:

- a. LDL-C Screening
- b. LDL-C Control (<100 mg/dL)

### **Non-HEDIS<sup>®</sup> Measure:**

#### **1. Statin Therapy for Persons with Cardiovascular Conditions**

Members with at least one statin prescription filled during the measurement year.

-----

## HYPERTENSION

### HEDIS® Measure:

#### 1. Controlling High Blood Pressure (CBP):

Percent members 18-85 years of age who had a diagnosis of HTN and whose BP was adequately controlled (<140/90 mm Hg) during the measurement year.

### Non-HEDIS® Measure:

#### 1. Percent Members with any Diagnosis of Hypertension

-----

## AC PHARYNGITIS

### HEDIS® Measure:

#### 1. Appropriate Testing for Children With Pharyngitis (CWP):

Children 2-18 years diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus test.

-----

## ADULT PREVENTIVE SERVICES (AGES 18 – 65+)

### HEDIS® Measures:

#### 1. Colorectal Cancer Screening (COL)

Percent members 50-80 years of age who had appropriate screening for colorectal cancer.

#### 2. Breast Cancer Screening (BCS)

Percent women 40-69 years of age who had a mammogram to screen for breast CA.

#### 3. Cervical Cancer Screening (CCS)

Percent women 21-64 years of age who received one or more Pap tests to screen for cervical cancer.

#### 4. Chlamydia Screening in Women (21-24 years) (CHL)

Percent women 21-24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.

#### 5. Flu Shots for Adults Ages 50-64 (CAHPS®) (FSA) [Commercial only]

A rolling average represents the percentage of commercial members 50-64 years of age who received an influenza vaccination between September 1 of the measurement year and the date on which the CAHPS® survey was completed.

#### 6. Flu Shots for Older Adults (CAHPS®) (FSO) [Medicare only]

Percent Medicare members 65 years of age and older as of January 1 of the measurement year who received an influenza vaccination between September 1 of the measurement year and the date on which the CAHPS® survey was completed.

#### 7. Pneumonia Vaccination Status for Older Adults (PNU) [Medicare only]

Percentage of Medicare members 65 years of age and older as of January 1 of the measurement year who have ever received a pneumococcal vaccine.

-----

## PREVENTIVE SERVICES FOR INFANTS AND CHILDREN (BIRTH – 24 MONTHS)

### HEDIS® Measures:

#### 1. **Childhood Immunization Status (CIS):**

Percent of children two years of age who had four DTaP, three IPV, one MMR, two Hib, three Hepatitis B, one VZV and four pneumococcal conjugate vaccines by their second birthday. The measure also calculates two separate combination rates, Combination 2 (DTaP, IPV, MMR, Hib, hepatitis B, VZV), and Combination 3 (DTaP, IPV, MMR, Hib, hepatitis B, VZV, pneumococcal conjugate).

#### 2. **Well-Child Visits in the First 15 Months of Life (W15):**

Percent members who turned 15 months old during the measurement year and who had six well-child visits with a PCP during their first 15 months of life.

#### 3. **Lead Screening in Children (LSC) [Medicaid only]**

Percent of children 2 years of age who had one or more capillary or venous lead blood tests for lead poisoning by their second birthday.

-----

## PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (AGES 2 - 21)

### HEDIS® Measures:

#### 1. **Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34):**

Percent members 3-6 years of age who received one or more well-child visits with a PCP during the measurement year.

#### 2. **Adolescent Well-Care Visits (AWC):**

Percent members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.

#### 3. **Chlamydia Screening in Women (16 – 20 years) (CHL):**

Percent of women 16-20 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

-----

## ROUTINE PRENATAL AND POSTNATAL CARE

### HEDIS® Measures:

#### 1. **Prenatal and Postpartum Care (PPC):**

- a. Prenatal Care: percent deliveries that received a prenatal care visit in the first trimester or within 42 days of enrollment.
  - b. Postpartum Care: percent deliveries that had a postpartum visit on or between 21 and 56 days after delivery.
- 

## CHRONIC KIDNEY DISEASE

### Non-HEDIS® Measure:

1. **Percent members in HEDIS® Diabetes and Hypertension populations who had a GFR test.**
-

## **MANAGEMENT OF ACUTE LOW BACK PAIN**

### **HEDIS® Measure:**

#### **1. Use of Imaging Studies for Low Back Pain (LBP):**

Percent members with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of diagnosis.

-----

## **MANAGEMENT OF UNCOMPLICATED ACUTE BRONCHITIS IN ADULTS**

### **HEDIS® Measure:**

#### **1. Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis (AAB):**

Percent of adults 18-64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.

-----

## **MQIC GUIDELINES THAT WILL NOT HAVE MEASURES AT THIS TIME:**

1. DVT
2. Heart Failure
3. Adult Obesity
4. Osteoarthritis
5. Adult Unintended Pregnancy Prevention
6. Childhood Overweight Prevention
7. Childhood Overweight Treatment
8. Osteoporosis