

2011 MQIC MEASURES SUMMARY LISTING FOR ANNUAL PERFORMANCE REPORTING

DIABETES

HEDIS® Measures:

1. **Comprehensive Diabetes Care (CDC):**

Percentage of members 18-75 years of age with diabetes (type 1 and type 2) who had each of the following:

- a. HbA1c Testing
- b. HbA1c Poor Control (>9.0%)
- c. HbA1c Control (<8.0%)
- d. HbA1c Good Control (<7.0%) (Additional exclusion criteria are required for this indicator)
- e. Eye Exam (retinal) performed
- f. LDL-C Screening
- g. LDL-C Control (<100 mg/dL)
- h. Medical Attention for Nephropathy
- i. Blood Pressure Control (<140/80 mm Hg)
- j. Blood Pressure Control (<140/90 mm Hg)

Non-HEDIS Measures:

1. **Statin Therapy for Persons with Diabetes**

Members with at least one statin prescription filled during the measurement year.

2. **Disease-Burden Measures** (reported every 5 years; due with HEDIS® 2013 data)

- a. Acute Myocardial Infarction (AMI)
- b. Cerebrovascular Accident (CVA)
- c. Amputations
- d. Dialysis

ASTHMA

HEDIS® Measures:

1. **Use of Appropriate Medications for People With Asthma (ASM):**

Percent members 5-50 years of age who were identified as having persistent asthma and who were appropriately prescribed medication during the measurement year.

- a. 5-11 Years Old
- b. 12-50 Years Old
- c. Total (sum of the two numerators divided by sum of the two denominators)

Non-HEDIS Measures:

1. **Periodic Assessment (5-50 Years Old)**

Members who had at least one preventive/ambulatory health services visit within the measurement year with a PCP, pulmonologist, or allergist.

2. **Emergency Department Visits per 1000 Members with Asthma and Being Treated for a Primary Diagnosis of Asthma**

TOBACCO CONTROL

HEDIS® Measures:

1. Medical Assistance With Smoking Cessation and Tobacco Use (MSC):

The following components assess different facets of providing medical assistance with smoking cessation:

- a. Advising Smokers and Tobacco Users to Quit
- b. Discussing Cessation Medications
- c. Discussing Cessation Strategies

Non-HEDIS Measure:

1. Percent Current Smokers 18 years and older

DEPRESSION

HEDIS® Measures:

1. Antidepressant Medication Management (AMM):

Percent members 18 years of age and older who were diagnosed with a new episode of major depression and treated with antidepressant medication, and who remained on an antidepressant medication treatment.

- a. Effective Acute Phase Treatment
 - b. Effective Continuation Phase Treatment
-

SUBSTANCE USE DISORDERS

HEDIS® Measures:

1. Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET):

Percent of adolescent and adult members with a new episode of alcohol or other drug (AOD) dependence who received the following:

- a. Initiation of AOD Treatment
 - b. Engagement of AOD Treatment
-

HYPERCHOLESTEROLEMIA

HEDIS® Measures:

1. Cholesterol Management for Patients with Cardiovascular Conditions (CMC):

Percent members 18-75 years of age who were discharged alive for AMI, CABG or PCI from January 1 to November 1 of the year prior to the measurement year, *or* who had a diagnosis of IVD during the measurement year and the year prior to the measurement year, who had each of the following during the measurement year:

- a. LDL-C Screening
- b. LDL-C Control (<100 mg/dL)

Non-HEDIS Measure:

1. Statin Therapy for Persons with Cardiovascular Conditions

Members with at least one statin prescription filled during the measurement year.

HYPERTENSION

HEDIS® Measure:

1. Controlling High Blood Pressure (CBP):

Percent members 18-85 years of age who had a diagnosis of HTN and whose BP was adequately controlled (<140/90 mm Hg) during the measurement year.

Non-HEDIS Measure:

1. Percent Members with any Diagnosis of Hypertension

AC PHARYNGITIS

HEDIS® Measure:

1. Appropriate Testing for Children With Pharyngitis (CWP):

Percent children 2-18 years diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus test for the episode.

ADULT PREVENTIVE SERVICES (AGES 18 – 65+)

HEDIS® Measures:

1. Colorectal Cancer Screening (COL)

Percent members 50-75 years of age who had appropriate screening for colorectal cancer.

2. Breast Cancer Screening (BCS)

Percent women 40-69 years of age who had a mammogram to screen for breast cancer.

3. Cervical Cancer Screening (CCS)

Percent women 21-64 years of age who received one or more Pap tests to screen for cervical cancer.

4. Chlamydia Screening in Women (21-24 years) (CHL)

Percent women 21-24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.

5. Flu Shots for Adults Ages 50-64 (CAHPS®) (FSA) [Commercial only]

A rolling average represents the percentage of commercial members 50-64 years of age who received an influenza vaccination between September 1 of the measurement year and the date on which the CAHPS® survey was completed.

6. Flu Shots for Older Adults (CAHPS®) (FSO) [Medicare only]

Percent Medicare members 65 years of age and older as of January 1 of the measurement year who received an influenza vaccination between September 1 of the measurement year and the date on which the CAHPS® survey was completed.

7. Pneumonia Vaccination Status for Older Adults (PNU) [Medicare only]

Percentage of Medicare members 65 years of age and older as of January 1 of the measurement year who have ever received a pneumococcal vaccine.

PREVENTIVE SERVICES FOR INFANTS AND CHILDREN (BIRTH – 24 MONTHS)

HEDIS® Measures:

1. Childhood Immunization Status (CIS):

Percent of children two years of age who had four DTaP, three IPV, one MMR, three Hib, three Hepatitis B, one VZV, four Pneumococcal conjugate, two Hepatitis A, two or three Rotavirus (number varies based on which vaccine is given), and two Influenza vaccines by their second birthday. The measure also calculates nine separate combination rates, see Table 1.

Table 1 – Combination Vaccinations

Combination	DTap	IPV	MMR	HiB	Hep B	VZV	PCV	Hep A	RV	Influenza
Combination 2*	x	x	x	x	x	x				
Combination 3*	x	x	x	x	x	x	x			
Combination 4	x	x	x	x	x	x	x	x		
Combination 5	x	x	x	x	x	x	x		x	
Combination 6	x	x	x	x	x	x	x			x
Combination 7	x	x	x	x	x	x	x	x	x	
Combination 8	x	x	x	x	x	x	x	x		x
Combination 9	x	x	x	x	x	x	x		x	x
Combination 10	x	x	x	x	x	x	x	x	x	x

*MQIC health plans report all CIS measures, including Combos 2-10, but only Combos 2 and 3 results are reported on the MQIC Annual Performance Measurement report.

2. Well-Child Visits in the First 15 Months of Life (W15):

Percent members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.

3. Lead Screening in Children (LSC) [Medicaid only]

Percent of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.

PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (AGES 2 - 21)

HEDIS® Measures:

1. Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34):

Percent members 3-6 years of age who received one or more well-child visits with a PCP during the measurement year.

2. Adolescent Well-Care Visits (AWC):

Percent members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.

3. Chlamydia Screening in Women (16 – 20 years) (CHL):

Percentage of women 16-20 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.

4. Immunizations for Adolescents (IMA):

Percent of adolescents 13 years of age who had one dose of meningococcal vaccine and one tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap) or one tetanus, diphtheria toxoids vaccine (Td) by their 13th birthday. This measure calculates a rate for each vaccine and one combination rate.

ROUTINE PRENATAL AND POSTNATAL CARE

HEDIS® Measures:

1. Prenatal and Postpartum Care (PPC):

- a. Timeliness of prenatal Care: percent deliveries that received a prenatal care visit as a member of the organization in the first trimester *or* within 42 days of enrollment in the organization.
 - b. Postpartum Care: percent deliveries that had a postpartum visit on or between 21 and 56 days after delivery.
-

CHRONIC KIDNEY DISEASE

Non-HEDIS Measure:

1. **Percent members in HEDIS® Diabetes or Hypertension populations who had a serum creatinine test (necessary for GFR)**
-

MANAGEMENT OF ACUTE LOW BACK PAIN

HEDIS® Measure:

1. Use of Imaging Studies for Low Back Pain (LBP):

Percent members with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of diagnosis.

MANAGEMENT OF UNCOMPLICATED ACUTE BRONCHITIS IN ADULTS

HEDIS® Measure:

1. Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis (AAB):

Percent of adults 18-64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.

ADULT OBESITY

HEDIS® Measure:

1. Adult BMI Assessment (ABA):

Percent members 18-74 years of age who had an outpatient visit and who had their body mass index (BMI) documented during the measurement year or the year prior the measurement year.

CHILDHOOD OBESITY

HEDIS® Measure:

1. Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC):

Percent members 3-17 years of age who had an outpatient visits with a PCP or OB/GYN and who had evidence of BMI percentile documentation, counseling for nutrition and counseling for physical activity during the measurement year. This measure evaluates whether BMI percentile is assessed rather than an absolute BMI value.

- a. 3-11 Years Old
- b. 12-17 Years Old
- c. Total (sum of the two age stratifications)

MQIC GUIDELINES THAT DO NOT HAVE MEASURES AT THIS TIME:

1. Deep Venous Thrombosis
2. Systolic Heart Failure
3. In Office Use of Sedation
4. Osteoarthritis
5. Osteoporosis
6. Adult Unintended Pregnancy Prevention
7. Prevention of Pregnancy in Adolescents 12-17 years